



POST OP INSTRUCTIONS FOR BIOPSY
Please read these instructions carefully

Sometimes the after-effects of oral surgery are quite minimal. If you have any questions regarding your surgery, please do not hesitate to call the number above. A 24-hour answering service is available to contact the doctor-on-call after hours. Calling during regular office hours will afford a faster response to your questions or concerns.

First Hour: Pain meds should be taken as soon as possible. When you get home take the gauze out and eat anything cold and soft (see note below) then take your pain meds. After eating, place new gauze over the surgical site(s) and obtain pressure over the surgical site(s) for another 20 minutes. Be sure to eat before taking pain medication (nausea can occur if taken on an empty stomach).

Gauze should be changed every 20-30 minutes to help stop the bleeding. Make sure you are biting down to keep the pressure on the wound. There is extra gauze in your bag. You can also use a tea bag. just moisten with warm water, squeeze the water out and place just like the gauze, changing every 20-30 minutes. Heavy bleeding should stop the day of surgery but can ooze into the next day. (This is normal)

Swelling: Use the ice pack to minimize any swelling on the outside of your face for 1 to 2 days. Place 20 minutes on, 20 minutes off, throughout the day and evening.

Diet: Your diet will consist of soft, cold or room temperature foods for 1-2 days. For example: jello, pudding, yogurt, ice cream etc. (anything you can cut with the side of a fork). Avoid anything hot or spicy. No hot tea, coffee, or soup as these can irritate the tissues. It is best to avoid foods like nuts, sunflower seeds, popcorn, etc., which may get lodged in the socket areas. Drink plenty of fluids. **DO NOT USE A STRAW FOR FIVE DAYS AFTER SURGERY**

Persistent Bleeding: This usually means that the gauze is being clenched between teeth only and you are not exerting pressure on the surgical area. Try repositioning the gauze, changing every 20-30 minutes. If bleeding remains uncontrolled, please call our office.

Sutures: If sutures were placed, they should dissolve on their own within 7-10 days. You would have been instructed if the doctor used a different kind of suture

Brushing and Rinses: You can brush your teeth today but avoid the surgical site. **After day three** you may use warm salt water to rinse if your wound is in the oral cavity. Use ¼ teaspoon of salt mixed with 8 oz. of warm water and gently swish as needed.

Activities: *If you had general anesthesia, once you get home, you should be assisted when walking. You should be mindful for several hours, please also be careful and stand slowly.*

We advise no activities for 1-2 days after your surgery. ***Absolutely** no driving of a vehicle for the first 24hrs if you had a general anesthesia. No strenuous exercise for 5 days. No swimming in a swimming pool or hot tub for 2 weeks.

If you have any questions regarding your surgery, please contact our office at **714-730-6767**, otherwise we will be contacting you later in the day to make sure you are doing well.